

Publikationer 2005

Bjerrum L, Gahrn-Hansen B, Munck AP. Praktiserende læger, der måler C-reaktivt protein, udskriver mindre antibiotika til patienter med sinusitis – sekundærpublikation. Ugeskr Læger 2005; 167: 2775-7.

Bjerrum L. Ufrivillig natlig vandladning kan behandles uden medicin. Nordisk Campbell Center Nr. 1 2005.

Bro F, Pedersen KM, Hansen DG, Kragstrup J, Munck AP. Hvem kommunikerer almen praksis med? Praktiserende lægers kontakter med social og sundhedsvæsenet. Ugeskr Læger 2005;167:2519-22.

Christensen PM, Brixen K, Kristiansen IS. Cost-effectiveness of alendronate in the prevention of osteoporotic fractures in Danish women. Basic & Clinical Pharmacology & Toxicology 2005; 96: 387-96.

Eriksson T, Maclure M, Kragstrup J. To what extent do mass media health messages trigger patients' contacts with their GPs? Br J Gen Pract 2005; 55: 212-7.

Green A, Støvring H, Andersen M, Beck-Nielsen H. The epidemic of type 2 diabetes is a statistical artefact. Diabetologica 2005 (netversion)

Halvorsen PA, Kristiansen IS. Decisions on drug therapies by numbers needed to treat: a randomized trial. Arch Intern Med 2005; 165 (10): 1140-6.

Hansen BL. Viden om tegn på kræft med henblik på tidlig diagnostik. Ugeskr Læger 2005; 167: 1280-4.

Jarlbæk L, Andersen M, Hallas J, Engholm G, Kragstrup J. Use of opioids in a Danish population-based cohort of cancer patients. J Pain Symptom Manage 2005; 29: 336-43.

Kristiansen IS, Bingefors C, Nyholm D, Isacson D. The cost-effectiveness of continuous duodenal delivery of levodopa (Duodopa®) in patients with severe Parkinson's disease (abstract). Movement Disorders 2005; 20: S80.

Kristiansen IS, Olsen JA. Foreldreløse legemidler og helseløse pasienter. Tidsskr Nor Lægeforen 2005; 125: 992.

Lofthus CM, Cappelen I, Osnes EK, Falch JA, Kristiansen IS, Medhus AW et al. Local and national electronic databases in Norway demonstrate a varying degree of validity. *J Clin Epidemiol* 2005; 58: 280-5.

Nexøe J, Kristiansen IS, Gyrd-Hansen D, Nielsen JB. Influence of number needed to treat, costs and outcome on preferences for a preventive drug. *Family Practice* 2005; 22: 126-31.

Nexøe J, Stauning JA, Aaen-Larsen B. Almen Praksis ved Universiteterne. *Ugeskr Læger* 2005; 167: 12.

Rasmussen HMS, Søndergaard J, Kampmann JP, Andersen M. General practitioners prefer prescribing indicators based on detailed information on individual patients: a Delphi study. *Eur J Clin Pharmacol* 2005; 61: 237-41.